

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12Q 1/68</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 98/59072</b> <b>(43) International Publication Date:</b> 30 December 1998 (30.12.98)
<b>(21) International Application Number:</b> PCT/US98/12779 <b>(22) International Filing Date:</b> 19 June 1998 (19.06.98) <b>(30) Priority Data:</b> 60/050,405      20 June 1997 (20.06.97)      US <b>(71) Applicant (for all designated States except US):</b> AFFYMETRIX, INC. [US/US]; 3380 Central Expressway, Santa Clara, CA 95051 (US). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> MATSUZAKI, Hajime [US/US]; 3380 Central Expressway, Santa Clara, CA 95051 (US). MURPHY, Eric, A. [US/US]; 3380 Central Express- way, Santa Clara, CA 95051 (US). <b>(74) Agents:</b> KAGAN, Sarah, A. et al.; Banner & Witcoff, Ltd., 11th floor, 1001 G Street, N.W., Washington, DC 20001-4597 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
<b>(54) Title:</b> METHODS AND COMPOSITIONS FOR MULTIPLEX AMPLIFICATION OF NUCLEIC ACIDS		
<b>(57) Abstract</b>  A method is described for predetermining ratios of primer pairs present in a single reaction vessel so as to achieve approximately equimolar yield of products. The ratios are determined as a function of the length of the amplicon and the length of other amplicons being simultaneously tested. The primers may desirably be for p53 gene sequences.		

***FOR THE PURPOSES OF INFORMATION ONLY***

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## METHODS AND COMPOSITIONS FOR MULTIPLEX AMPLIFICATION OF NUCLEIC ACIDS

### RELATED APPLICATION

This application claims priority to U.S. Provisional Application, Serial No.  
5 60/050,405, filed on June 20, 1997, the text of which is expressly incorporated  
herein.

### BACKGROUND OF THE INVENTION

The polymerase chain reaction (PCR) is a simple and versatile method to  
amplify *in vitro* a specific segment of DNA for subsequent study (Saiki *et al.*,  
10 *Science* 230:1350 (1985); Saiki *et al.*, *Science* 235:487 (1985)). The PCR method  
has gained widespread use in biomedical research, and has revolutionized the  
accurate and early diagnosis of many inherited and acquired genetic disorders  
(Eisenstein, N. *Engl. J. Med.* 322:178 (1990)), particularly those caused by point  
mutations or small insertions or deletions including sickle cell anemia (Saiki *et al.*,  
15 *Science* 230:1350 (1985)), hemophilia A (Kogan *et al.*, *N. Engl. J. Med.* 317:985  
(1987)), Tay-Sach's disease (Myerowitz, *Proc. Natl. Acad. Sci. USA* 85:3955  
(1988); Myerowitz *et al.*, *J. Biol. Chem.* 263:18587 (1988)), cystic fibrosis  
(Riordan *et al.*, *Science* 245:1066 (1989)), and many others. With PCR, it is also  
possible to detect heterozygotic carriers in recessive disorders.

Polymerase chain reaction (PCR) is used for a variety of purposes. PCR can be used to amplify genomic DNA or other sources of nucleic acids for analysis. It is often desirable to be able to achieve equimolar yields of different length amplicons when performing multiplex PCR or multiple PCR reactions. Having an

5 approximately equimolar yield of amplicons is particularly useful, for example, when approximately equal representation of certain regions of genomic DNA amplified after multiplex PCR is desired. Prior to the methods of present invention, finding the appropriate experimental conditions useful to achieve this result has been difficult because PCR amplifies nucleic acids having different lengths with different

10 efficiencies. The yield of longer amplicons is often less than the yield of shorter amplicons because of those differences in PCR amplification efficiency. Figure 1 shows the difference in yields that one might expect, for example, when starting with equal primer concentrations used to amplify amplicons of varying lengths: A, B, C. There is a continuing need in the art for methods which permit the amplification of

15 different sequences with the same efficiency so that approximately equimolar products result.

#### **SUMMARY OF THE INVENTION**

It is an object of the present invention to provide a method of performing multiplex PCR which achieve approximately equimolar products.

20 It is another object of the invention to provide a set of primers for amplification of p53.

It is yet another object of the invention to provide a set of primers for amplification of p53 to achieve approximately equimolar products.

It is still another object of the invention to provide a mixture of primers for

performing multiplex PCR.

These and other objects of the invention are provided by one or more of the embodiments provided below. In one embodiment of the invention a method of performing multiple polymerase chain reactions in a single vessel is provided. The method comprises the steps of priming DNA synthesis on a template in a vessel with at least two sets of primers. The primers are present in the vessel at a predetermined ratio which is described by the formula:

$$C_A = C_L (L_A \div L_L)^2$$

$C_A$  is the concentration of primers for an amplicon A.  $C_L$  is the concentration of primer for the longest amplicon.  $L_A$  is the length of the amplicon A.  $L_L$  is the length of the longest amplicon.

Another embodiment provided by the invention is a method of performing multiple polymerase chain reactions in a single vessel. The method comprises priming DNA synthesis on a genomic p53 template in a vessel with ten sets of primers which amplify exons 2-11 of p53. The primers are shown in SEQ ID NO: 1-20. The primers are present in the vessel at the following ratio: exon 2 (89.4): exon 3 (26.9): exon 4 (450): exon 5 (245.8): exon 6 (138.3): exon 7 (101.8): exon 8 (193.0): exon 9 (70.8): exon 10 (146.5): exon 11 (177.3).

According to still another embodiment of the invention a set of primers for performing multiple polymerase chain reactions in a single vessel is provided. The set comprises twenty primers having sequences as shown in SEQ ID NO: 1-20.

According to yet another embodiment of the invention a mixture of primers for performing multiplex polymerase chain reaction is provided. The primers are present in the mixture at a predetermined ratio to each other. The ratio of the concentrations of

the primers is described by:

$$C_A = C_L (L_A \div L_L)^2$$

wherein  $C_A$  is the concentration of primers for an amplicon A; wherein  $C_L$  is the concentration of primer for the longest amplicon; wherein  $L_A$  is the length of the amplicon A; and wherein  $L_L$  is the length of the longest amplicon.

The present invention thus provides the art with a method useful for performing multiplex PCR. This method is particularly useful for amplification of multiple exons of p53. Moreover, a particular primer set useful for performing such multiplex PCR is also provided.

#### **BRIEF DESCRIPTION OF THE DRAWINGS**

Figure 1 shows the difference in yields that one might expect, for example, when starting with equal primer concentrations used to amplify amplicons of varying lengths: A, B, C.

Figure 2 illustrates the relationship for given values X and  $L_L$ , using the amplicons from different exons of the human p53 gene as an example.

#### **DETAILED DESCRIPTION OF THE INVENTION**

It is a discovery of the present invention that approximately equimolar yields of amplicons of varying lengths can be easily produced by multiplex PCR. It has been determined that varying the primer concentrations as a function of the lengths of amplicons yields approximately equimolar amounts of amplicons of varying lengths. The relationship between primer concentration and the length of amplicons is as follows:

$$C_A = C_L (L_A / L_L)^X$$

wherein  $C_A$  = the concentration of primers for an amplicon A;

$C_L$  = the concentration of primer for the longest amplicon;

$L_A$  = the length of amplicon A;

$L_L$  = the length of the longest amplicon; and

5 X is usually not zero and is often between one and three.

This relationship can be placed in a computer readable medium or be used with a computer system if desired.

Figure 2 illustrates the relationship for given values X and  $L_L$ , using the amplicons from different exons of the human p53 gene as an example. Using primer concentrations as set forth, for example in Figure 2, one skilled in the art can determine the optimum set of primer concentrations to yield approximately equimolar yields of varying length amplicons in a multiplex or multiple PCR. Preferably, primers having both comparable base composition and comparable melting temperatures are used. Also preferably,  $Mg^{+2}$  concentration, annealing temperatures, and cycling times of the PCR are optimized prior to choosing the desired set of primer concentrations in accordance with the present invention.

10

15

PCR techniques applicable to the present invention include *inter alia* those discussed in PCR PRIMER: A LABORATORY MANUAL, Dieffenbach, C.W. and Dveksler, G.S., eds., Cold Spring Harbor Laboratory Press (1995).

20 The present application further provides primer sequences, primer concentrations, and experimental conditions useful in the amplification of the coding region of the human p53 gene. Particularly useful primers for amplification of exons of the p53 gene are set forth in Table 1.



TABLE 1

## p53 Primer Set

20 primers in 1 mM Tris-HCl, pH 7.4, 0.1 mM EDTA, sequences:

5	Exon 2:	5'-TCATGCTGGATCCCCACTTTTCCTCTTG-3'
		5'TGGCCTGCCCTTCCAATGGATCCACTCA-3'
10	Exon 3:	5'-AATTCATGGGACTGACTTTCTGCTCTTGTC-3'
		5'-TCCAGGTCCCAGCCCAACCCTTGTCC-3'
15	Exon 4:	5'-GTCCTCTGACTGCTCTTTTCACCCATCTAC-3'
		5'-GGGATACGGCCAGGCATTGAAGTCTC-3'
20	Exon 5:	5'-CTTGTGCCCTGACTTTCAACTCTGTCTC-3'
		5'-TGGGCAACCAGCCCTGTCGTCTCTCCA-3'
	Exon 6:	5'-CCAGGCCTCTGATTCCTCACTGATTGCTC-3'
		5'-GCCACTGACAACCACCCTTAACCCCTC-3'
	Exon 7:	5'-GCCTCATCTTGGGCCTGTGTTATCTCC-3'
		5'-GGCCAGTGTGCAGGGTGGCAAGTGGCTC-3'
	Exon 8:	5'-GTAGGACCTGATTTCTTACTGCCTCTTGC-3'
		5'-ATAACTGCACCCTTGGTCTCCTCCACCGC-3'
	Exon 9:	5'-CACTTTTATCACCTTTCCTTGCCTCTTTCC-3'
		5'-AACTTTCCACTTGATAAGAGGTCCCAAGAC-3'
	Exon 10:	5'-ACTTACTTCTCCCCCTCCTCTGTTGCTGC-3'
		5'-ATGGAATCCTATGGCTTTCCAACCTAGGAAG-3'
	Exon 11:	5'-CATCTCTCCTCCCTGCTTCTGTCTCCTAC-3'
		5'-CTGACGCACACCTATTGCAAGCAAGGGTTC-3'



Table 2 shows particularly useful concentrations of the primers set forth in Table 1 for multiplex PCR amplification using the experimental conditions set forth in Table 3.

TABLE 2

## Primer Concentrations in p53 Primer Set

Values of X Typical values of  $C_L$  

	Amplicon	Length	Primer Concs
Longest	4	368 bp	450.0 nM
	5	272 bp	245.8 nM
	8	241 bp	193.0 nM
	11	231 bp	177.3 nM
	10	210 bp	146.5 nM
	6	204 bp	138.3 nM
	7	175 bp	101.8 nM
	2	164 bp	89.4 nM
	9	146 bp	70.8 nM
Shortest	3	90 bp	26.9 nM

TABLE 3

**Multiplex PCR**

Start with 250 ng of Template DNA.

PCR Components for 100 ul PCR in 0.2 ml thin walled tubes:

	Stock Conc	Final Conc	for 1 reaction
Buffer (No Mg)	10 X	1 X	10.0 ul
MgCl <sub>2</sub>	25 mM	2.5 mM	10.0 ul
dATP	10 mM	200 uM	2.0 ul
dCTP	10 mM	200 uM	2.0 ul
dGTP	10 mM	200 uM	2.0 ul
dTTP	10 mM	200 uM	2.0 ul
Taq GOLD	5 U/ul	10 U	2.0 ul
p53 Primer Set	20 X	1 X	5.0 ul
Water			
Human genomic DNA		250 ng	
Total Volume			100.0 ul

Final Concentrations in Buffer (No Mg) are 10 mM Tris-HCl (pH 8.3), 50 mM KCl  
Taq GOLD is AmpliTaq Gold™ from Perkin Elmer catalog # N808-0243

**PCR Cycles:**

	94 C	10 min
35 Cycles:	94 C	30 sec
	60 C	30 sec
	72 C	45 sec
	72 C	10 min

**To visualize amplicons by gel Analysis:**

Visualize PCR products on 4% NuSieve Agarose Gel

NuSieve™ Agarose 3:1 is from FMC catalog # 50092

Load 5 ul of PCR + loading buffer

Use 50 bp Ladder (Gibco/BRL catalog # 10416-014) as size marker

Run gel at 125 Volts for 30 min. to 90 min.

**Expected PCR Products:**

Amplicon	Length
Exon 2	164 bp
Exon 3	90 bp
Exon 4	368 bp
Exon 5	272 bp
Exon 6	204 bp
Exon 7	175 bp
Exon 8	241 bp
Exon 9	146 bp
Exon 10	210 bp
Exon 11	225 bp

**Order in Gel:**

Amplicon	Length
Exon 4	368 bp
Exon 5	272 bp
Exon 8	241 bp
Exon 11	225 bp
Exon 10	210 bp
Exon 6	204 bp
Exon 7	175 bp
Exon 2	164 bp
Exon 9	146 bp
Exon 3	90 bp

Using the methods and reagents provided herein, we achieved multiplex PCR amplification of coding regions shown of the human p53 gene in approximately equimolar amounts. That desirable result was achieved in a single-tube reaction. The achievement of such desirable results with the remarkable convenience of a single tube reaction further illustrates the contribution to the art made by the present invention.

The methods and compositions of the present invention are useful in virtually any context in which equimolar yields of various PCR products are desired. Such contexts include without limitation paternity testing, forensic analysis, genetic screening, polymorphism detection, and mutation analyses. The present invention can be used to amplify nucleic acids for all forms of sequence analysis known to those skilled in the art. Sequence analysis techniques includes, for example, dideoxy-sequencing and sequence analysis using high-density nucleic acid arrays: the GeneChip® probe arrays or VLSIPS™ technology of Affymetrix, Inc. High density nucleic acid arrays are discussed for example in Chee, M., Yang, R., Hubbell, E., Berno, A., Huang, X.C., Stern, D., Winkler, J., Lockhart, D.J., Morris, M.S., & Fodor, S.P., *Science* **5287**, 610-614 (1996), U.S. Patent No. 5,445,934, and International Publication No. WO 95/11995 corresponding to PCT Application No. PCT/US94/12305.

The p53 gene and its protein product are discussed in *Molecular Biology of the Cell*, 3rd Edition, Alberts, B., Bray, D., Lewis, J., Raff, M., Roberts, K., and Watson, J.D., Garland Publishing (1994) at pages 889 and 1284-1289.

It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of any appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety for all purposes.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION

(i) APPLICANT: MATSUZAKI, HAJIME

(ii) TITLE OF THE INVENTION: METHODS AND COMPOSITIONS FOR  
MULTIPLEX AMPLIFICATION OF NUCLEIC ACIDS

(iii) NUMBER OF SEQUENCES: 20

## (iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Banner & Witcoff  
(B) STREET: 1001 G Street, NW  
(C) CITY: Washington  
(D) STATE: DC  
(E) COUNTRY: USA  
(F) ZIP: 20001

## (v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette  
(B) COMPUTER: IBM Compatible  
(C) OPERATING SYSTEM: DOS  
(D) SOFTWARE: FastSEQ for Windows Version 2.0

## (vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:  
(B) FILING DATE: 28-JUN-1998  
(C) CLASSIFICATION:

## (vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 60/050,405  
(B) FILING DATE: 20-JUN-1997

## (viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Kagan, Sarah A  
(B) REGISTRATION NUMBER: 32141  
(C) REFERENCE/DOCKET NUMBER: 03848.74891

## (ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 202-508-9100  
(B) TELEFAX: 202-508-9299  
(C) TELEX:

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 28 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TCATGCTGGA TCCCCACTTT TCCTCTTG  
28

5

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

10

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

TGGCCTGCCC TTCCAATGGA TCCACTCA  
28

(2) INFORMATION FOR SEQ ID NO:3:

15

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

20

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

AATTCATGGG ACTGACTTTC TGCTCTTGTC  
30

(2) INFORMATION FOR SEQ ID NO:4:

25

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

30

TCCAGGTCCC AGCCCAACCC TTGTCC  
26

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

35

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

GTCTCTGAC TGCTCTTTC ACCCATCTAC  
30

(2) INFORMATION FOR SEQ ID NO:6:

5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 26 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

GGGATACGGC CAGGCATTGA AGTCTC  
26

(2) INFORMATION FOR SEQ ID NO:7:

15 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

20 CTTGTGCCCT GACTTTC AAC TCTGTCTC  
28

(2) INFORMATION FOR SEQ ID NO:8:

25 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

30 TGGGCAACCA GCCCTGTCGT CTCTCCA  
27

(2) INFORMATION FOR SEQ ID NO:9:

35 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 29 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CCAGGCCTCT GATTCCTCAC TGATTGCTC  
29

(2) INFORMATION FOR SEQ ID NO:10:

5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GCCACTGACA ACCACCCTTA ACCCCTC  
27

(2) INFORMATION FOR SEQ ID NO:11:

15 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 27 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

20 GCCTCATCTT GGGCCTGTGT TATCTCC  
27

(2) INFORMATION FOR SEQ ID NO:12:

25 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 28 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

30 GGCCAGTGTG CAGGGTGGCA AGTGGCTC  
28

(2) INFORMATION FOR SEQ ID NO:13:

35 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 30 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:



GTAGGACCTG ATTCCTTAC TGCCTCTTGC  
30

(2) INFORMATION FOR SEQ ID NO:14:

- 5 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 29 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

10 ATAACTGCAC CCTTGGTCTC CTCCACCGC  
29

(2) INFORMATION FOR SEQ ID NO:15:

- 15 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 30 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

20 CACTTTTATC ACCTTTCCTT GCCTCTTTCC  
30

(2) INFORMATION FOR SEQ ID NO:16:

- 25 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 30 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

AACTTTCCAC TTGATAAGAG GTCCCAAGAC  
30

30 (2) INFORMATION FOR SEQ ID NO:17:

- 35 (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 29 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

ACTTACTTCT CCCCCTCCTC TGTTGCTGC

29

## (2) INFORMATION FOR SEQ ID NO:18:

## (i) SEQUENCE CHARACTERISTICS:

- 5 (A) LENGTH: 31 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

10 ATGGAATCCT ATGGCTTTCC AACCTAGGAA G  
31

## (2) INFORMATION FOR SEQ ID NO:19:

## (i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 29 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

20 CATCTCTCCT CCCTGCTTCT GTCTCCTAC  
29

20

## (2) INFORMATION FOR SEQ ID NO:20:

## (i) SEQUENCE CHARACTERISTICS:

- 25 (A) LENGTH: 30 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

CTGACGCACA CCTATTGCAA GCAAGGGTTC  
30

**Claims**

1. A method of performing multiple polymerase chain reactions in a single vessel, comprising:

priming DNA synthesis on a template in a vessel with at least two sets  
 5 of primers, wherein the primers are present in the vessel at a predetermined ratio, wherein the ratio is described by:

$$C_A = C_L (L_A \div L_L)^2$$

wherein  $C_A$  is the concentration of primers for an amplicon A; wherein  $C_L$  is the concentration of primer for the longest amplicon; wherein  $L_A$  is the length of the  
 10 amplicon A; and wherein  $L_L$  is the length of the longest amplicon.

2. The method of claim 1 wherein the template is genomic DNA encoding p53.
3. The method of claim 1 wherein the template is a cDNA encoding p53.
4. The method of claim 1 wherein the primers amplify at least 2 exons of p53 selected from the group consisting of exons 2-11.
- 15 5. The method of claim 1 wherein the primers amplify at least 4 exons of p53 selected from the group consisting of exons 2-11.
6. The method of claim 1 wherein the primers amplify exons 2-11 of p53.
7. The method of claim 4 wherein the primers are selected from those shown in SEQ ID NO: 1-20.
- 20 8. The method of claim 5 wherein the primers are selected from those shown in SEQ ID NO: 1-20.
9. The method of claim 6 wherein the primers are shown in SEQ ID NO: 1-20.
10. The method of claim 9 wherein the primers are present in the following ratios:  
 25 exon 2 (89.4): exon 3 (26.9): exon 4 (450): exon 5 (245.8): exon 6 (138.3):  
 exon 7 (101.8): exon 8 (193.0): exon 9 (70.8): exon 10 (146.5): exon 11 (177.3).
11. A method of performing multiple polymerase chain reactions in a single vessel, comprising:

30 priming DNA synthesis on a genomic p53 template in a vessel with ten

sets of primers which amplify exons 2-11 of p53, wherein the primers are shown in SEQ ID NOS: 1-20, wherein the primers are present in the vessel at the following ratios: exon 2 (89.4), exon 3 (26.9), exon 4 (450), exon 5 (245.8), exon 6 (138.3), exon 7 (101.8), exon 8 (193.0), exon 9 (70.8), exon 10 (146.5), exon 11 (177.3).

- 5 12. A kit comprising a set of primers for performing multiple polymerase chain reactions in a single vessel, comprising:

twenty primers having sequences as shown in SEQ ID NO: ID NOS: 1-20.

- 10 13. The kit of claim 12 wherein the ratio of the concentrations of the primers is described by:

$$C_A = C_L (L_A \div L_L)^2$$

wherein  $C_A$  is the concentration of primers for an amplicon A; wherein  $C_L$  is the concentration of primer for the longest amplicon; wherein  $L_A$  is the length of the amplicon A; and wherein  $L_L$  is the length of the longest amplicon.

- 15 14. The kit of claim 12 wherein the ratio of the primers is: exon 2 (89.4): exon 3 (26.9): exon 4 (450): exon 5 (245.8): exon 6 (138.3): exon 7 (101.8): exon 8 (193.0): exon 9 (70.8): exon 10 (146.5): exon 11 (177.3).

- 20 15. A mixture of primers for performing multiplex polymerase chain reaction, wherein the primers are present in the mixture at a predetermined ratio to each other, wherein the ratio of the concentrations of the primers is described by:

$$C_A = C_L (L_A \div L_L)^2$$

wherein  $C_A$  is the concentration of primers for an amplicon A; wherein  $C_L$  is the concentration of primer for the longest amplicon; wherein  $L_A$  is the length of the amplicon A; and wherein  $L_L$  is the length of the longest amplicon.

- 25 16. The mixture of claim 15 which comprises at least 4 primers.  
 17. The mixture of claim 15 which comprises at least 6 primers.  
 18. The mixture of claim 15 which comprises at least 8 primers.  
 19. The mixture of claim 15 which comprises at least 10 primers.  
 20. The mixture of claim 15 which comprises at least 12 primers.  
 30 21. The mixture of claim 15 which comprises at least 14 primers.  
 22. The mixture of claim 15 which comprises at least 16 primers.

23. The mixture of claim 15 which comprises at least 18 primers.
24. The mixture of claim 15 which comprises at least 20 primers.





FIG. 2

MULTIPLEX PCR PRIMER CONCENTRATIONS AS A FUNCTION OF THE LENGTHS OF AMPLICONS

VALUES OF X		2	2	2	1	0	3	
TYPICAL VALUES OF L <sub>L</sub>		1000 nM	500 nM	250 nM	500 nM	500 nM	500 nM	
	AMPLI- CON	LENGTH	X=2, L <sub>L</sub> =1000	X=2, L <sub>L</sub> =500	X=2, L <sub>L</sub> =250	X=1, L <sub>L</sub> =500	X=0, L <sub>L</sub> =500	X=3, L <sub>L</sub> =500
LONGEST	4	368 bp	1000.0 nM	500.0 nM	250.0 nM	500.0 nM	500.0 nM	500.0 nM
	5	272 bp	546.3 nM	273.2 nM	136.6 nM	369.6 nM	500.0 nM	201.9 nM
	8	241 bp	428.9 nM	214.4 nM	107.2 nM	327.4 nM	500.0 nM	140.4 nM
	11	231 bp	394.0 nM	197.0 nM	98.5 nM	313.9 nM	500.0 nM	123.7 nM
	10	210 bp	325.6 nM	162.8 nM	81.4 nM	285.3 nM	500.0 nM	92.9 nM
	6	204 bp	307.3 nM	153.7 nM	76.8 nM	277.2 nM	500.0 nM	85.2 nM
	7	175 bp	226.1 nM	113.1 nM	56.5 nM	237.8 nM	500.0 nM	53.8 nM
	2	164 bp	198.6 nM	99.3 nM	49.7 nM	222.8 nM	500.0 nM	44.3 nM
SHORTEST	9	146 bp	157.4 nM	78.7 nM	39.4 nM	198.4 nM	500.0 nM	31.2 nM
	3	90 bp	59.8 nM	29.9 nM	15.0 nM	122.3 nM	500.0 nM	7.3 nM

MULTIPLEX [PRIMERS] AS A FUNCTION OF THE LENGTHS OF AMPLICONS





# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 98/12779

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 96 01909 A (VISIBLE GENETICS INC ;DIAMANDIS ELEFThERIOS (CA); DUNN JAMES M (CA) 25 January 1996 see page 11, line 1 - page 14, line 6 see page 19 - page 29 see claims 16-33 ---	1-24
A	WO 96 10648 A (PROMEGA CORP) 11 April 1996 see page 9, line 23 - page 35 ---	1-24
A	WO 96 39535 A (VIJG JAN ;LI DAIZONG (US)) 12 December 1996 see page 7, line 15 - page 10, line 14 ---	1-24
A	EP 0 648 845 A (EASTMAN KODAK CO) 19 April 1995 see page 2, line 49 - page 3, line 2 see page 5, line 37 - line 47 ---	1-24
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

28 October 1998

Date of mailing of the international search report

06/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Osborne, H

# INTERNATIONAL SEARCH REPORT

Int'l Application No  
PCT/US 98/12779

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WO 96 41012 A (GENZYME CORP ; SHUBER ANTHONY P (US)) 19 December 1996 see the whole document -----</p>	1

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/12779

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9601909 A	25-01-1996	US 5545527 A US 5552283 A AU 3094895 A AU 690795 B AU 4321696 A CA 2191233 A EP 0770145 A EP 0760011 A JP 10504897 T WO 9607761 A	13-08-1996 03-09-1996 09-02-1996 30-04-1998 27-03-1996 14-03-1996 02-05-1997 05-03-1997 12-05-1998 14-03-1996
WO 9610648 A	11-04-1996	AU 3998195 A CA 2118048 A	26-04-1996 31-03-1996
WO 9639535 A	12-12-1996	AU 5775596 A CA 2223061 A EP 0873419 A US 5814491 A	24-12-1996 12-12-1996 28-10-1998 29-09-1998
EP 0648845 A	19-04-1995	JP 7163370 A US 5674717 A	27-06-1995 07-10-1997
WO 9641012 A	19-12-1996	AU 6163696 A CA 2223729 A EP 0832290 A	30-12-1996 19-12-1996 01-04-1998